

Uniform pathwise connectedness and Whitney levels

Iwona Krzemińska

Opole Technical University, Department of Mathematics, ul. Luboszycka 3, 45-036 Opole, Poland

Received 9 September 1998; received in revised form 2 July 1999

Abstract

It is proved that the property of being a uniformly pathwise connected continuum is a Whitney property, but is not a Whitney-reversible property. © 2001 Elsevier Science B.V. All rights reserved.

Keywords: Continuum; Hyperspace of subcontinua; Whitney property; Uniform pathwise connectedness

AMS classification: 54B20; 54F15

Let X be a continuum, i.e., a compact connected metric space, and let $C(X)$ be the space of all nonempty subcontinua of X topologized by the Hausdorff metric.

A continuum X is said to be *uniformly pathwise connected* if there exists a family $\mathcal{F} = \{p : [a, b] \rightarrow X\}$ of paths in X , satisfying

- (*) for any $x, y \in X$ there is a path $p \in \mathcal{F}$ joining x with y , and
- (**) for any $\varepsilon > 0$ there is a positive integer n such that for each $p \in \mathcal{F}$ there are numbers $t_0 = a < t_1 < t_2 < \dots < t_n = b$ such that $\text{diam } p([t_{i-1}, t_i]) \leq \varepsilon$ for $i = 1, 2, \dots, n$.

It is an easy observation that the continuous images of the cone over the Cantor set are uniformly pathwise connected. The converse is also true and was proved by Kuperberg [8].

A map $\mu : C(X) \rightarrow [0, \infty)$ satisfying the conditions

$$\mu \text{ is continuous on } C(X), \quad (1)$$

$$\text{if } A \subset B \text{ and } A \neq B, \text{ then } \mu(A) < \mu(B), \quad (2)$$

$$\mu(\{x\}) = 0 \text{ for every } x \in X, \quad (3)$$

is called a *Whitney map*.

E-mail address: ikrze@polo.po.opole.pl (I. Krzemińska).

The Whitney maps are open and monotone (see Nadler [10, p. 400]). Then it is natural to study the following properties.

A property \mathcal{P} is said to be a *Whitney property* provided that whenever a continuum X has property \mathcal{P} , so does $\mu^{-1}(t)$ for each Whitney map μ , for $C(X)$ and each t , $0 \leq t < \mu(X)$ (see [10, p. 399]).

A property \mathcal{P} is said to be a *Whitney-reversible property* provided that whenever X is a continuum such that $\mu^{-1}(t)$ has property \mathcal{P} for all Whitney maps μ for $C(X)$ and all $0 < t < \mu(X)$, then X has property \mathcal{P} (see [10, p. 453]).

It is known that arcwise connectedness is a Whitney property (see [10, p. 405]). In [5] Kato has proved that if X is uniformly pathwise connected continuum and has the property of Kelley (see [10, p. 538] for the definition), then for any Whitney map μ , and any $t \in [0, \mu(X))$, the level $\mu^{-1}(t)$ is uniformly pathwise connected. In this paper we prove that the same conclusion holds without the assumption that X has the property of Kelley. This answers in the positive the question posed by Illanes and Nadler in [4]. There are easy examples of uniformly pathwise connected continua without the property of Kelley.

It is known that arcwise connectedness is not a Whitney-reversible property (see [10, p. 457]). Illanes and Nadler asked in the book [4] if uniform pathwise connectedness is a Whitney-reversible property. In this paper we show that the assumption that all Whitney levels $\mu^{-1}(t)$, for $0 < t < \mu(X)$, are uniformly pathwise connected does not suffice for the level $\mu^{-1}(0)$ to have this property.

An *order arc* in $C(X)$ is an arc \mathcal{A} in $C(X)$ such that for every pair of elements L, K of \mathcal{A} , we have $L \subset K$ or $K \subset L$. If A and B are subcontinua of X , and $A \subset B$, then there is an order arc in $C(X)$ beginning at A and ending at B (see [10, p. 63], where detailed references can be found).

By a path in X joining points $x, y \in X$ we mean a continuous function $p: [a, b] \rightarrow X$ such that $p(a) = x$ and $p(b) = y$.

The symbol $N_\delta(A)$ denotes the open δ -neighborhood of A in X ; dist denotes the Hausdorff distance.

We will use the following lemma (cf. Krasinkiewicz [9, p. 157]).

Lemma 1. *For any $\varepsilon > 0$ there exists $\delta > 0$ such that if K and L are elements of $C(X)$ such that $K \subset N_\delta(L)$ and $|\mu(L) - \mu(K)| < \delta$, then $\text{dist}(K, L) < \varepsilon$.*

Modifying the proof of (14.8.1) from [10, p. 405], we get the following.

Lemma 2. *Let μ be a fixed Whitney map and let $t \in (0, \mu(X))$. There exists a family \mathcal{F} of paths $q: [0, t] \rightarrow \mu^{-1}(t)$ such that for any $\varepsilon > 0$ there exists a positive integer n such that if $A, B \in \mu^{-1}(t)$ and $A \cap B \neq \emptyset$ then we can choose a path $q \in \mathcal{F}$ such that $q(0) = B$ and $q(t) = A$ and such that the diameter of $q([(i-1)\frac{t}{n}, i\frac{t}{n}])$ is $\leq \varepsilon$, for $i = 1, \dots, n$.*

Proof. If $A, B \in \mu^{-1}(t)$ and $A \cap B \neq \emptyset$ then for $x \in A \cap B$ consider order arcs \mathcal{A} and \mathcal{B} in $C(X)$ one with ends $\{x\}$ and A and other with ends $\{x\}$ and B . Let α and

β be parametrizations of these arcs by using the Whitney map, i.e., $\alpha: [0, t] \rightarrow \mathcal{A}$, and $\beta: [0, t] \rightarrow \mathcal{B}$, such that

$$\mu(\alpha(u)) = \mu(\beta(u)) = u, \quad (4)$$

for each $u \in [0, t]$. We have $\alpha(0) = \beta(0) = \{x\}$, and $\alpha(t) = A$ and $\beta(t) = B$.

For a fixed $u \in [0, t]$ the subset $\{\alpha(u) \cup \beta(w): w \in [0, t]\}$, of $C(X)$ is connected. Thus, from the continuity of μ it easily follows that for any $u \in [0, t]$ there exists $s \in [0, t]$ such that

$$\mu(\alpha(u) \cup \beta(s)) = t.$$

Observe that the set $T(u) = \{s: \mu(\alpha(u) \cup \beta(s)) = t\}$ is a closed subset of $[0, t]$. Thus, $s(u) = \sup T(u)$ belongs to $T(u)$ and therefore we can define a function $q: [0, t] \rightarrow \mu^{-1}(t)$, setting

$$q(u) = \alpha(u) \cup \beta(s(u)), \quad \text{for } u \in [0, t]. \quad (5)$$

Although $s(u)$ need not be continuous, the function q is continuous.

Indeed, let $u_0 \in [0, t]$ and let u_n , where $n = 1, 2, \dots$, be a sequence of points of $[0, t]$ convergent to u_0 . Let us denote by v arbitrarily chosen point of condensation of the set $\{s(u_n): n = 1, 2, \dots\}$. The sequence $s(u_n)$ has a subsequence $s(u_{n_k})$ convergent to v . Thus, the sequence $q(u_{n_k})$ converges to $\alpha(u_0) \cup \beta(v)$, and $\alpha(u_0) \cup \beta(v) \in \mu^{-1}(t)$, for compactness of the level $\mu^{-1}(t)$.

Thus, to prove the continuity of q , it suffices to show that

$$\alpha(u_0) \cup \beta(v) = \alpha(u_0) \cup \beta(s(u_0)), \quad (6)$$

as v has been taken arbitrarily.

To do this, observe that $v \leq s(u_0)$, for $v \in T(u_0)$. Thus, $\alpha(u_0) \cup \beta(v) \subset \alpha(u_0) \cup \beta(s(u_0))$. Since the sets on both sides of this inclusion lie on $\mu^{-1}(t)$, hence, using (2) from the definition of the Whitney map, we obtain the equality (6).

Now, fix $\varepsilon > 0$ and take $\delta > 0$ according to Lemma 1. Then, for this δ take $\delta' > 0$ according to Lemma 1. We have $\text{dist}(K, L) < \delta$ for $K, L \in C(X)$ whenever $K \subset N_{\delta'}(L)$ and $|\mu(K) - \mu(L)| < \delta'$.

Let the integer n be such that $\frac{t}{n} < \delta'$.

To show that the number n is that one which we need for the function q , let u and w , where $u < w$, be points of $[0, t]$ such that

$$u \text{ and } w \text{ belong to } \left[(i-1)\frac{t}{n}, i\frac{t}{n}\right], \quad \text{for some } i \in \{1, \dots, n\}. \quad (7)$$

Since $u < w$, then

$$\alpha(u) \subset \alpha(w) \quad (8)$$

and, by the monotonicity of the function s , we have $\beta(s(w)) \subset \beta(s(u))$.

Since $w - u < \delta'$ (see (7)), then, by (4), $\mu(\alpha(w)) - \mu(\alpha(u)) < \delta'$. Thus, by (8) and Lemma 1, we have $\text{dist}(\alpha(u), \alpha(w)) < \delta$.

Therefore, using inclusions

$$\alpha(w) \subset N_{\delta}(\alpha(u)) \quad \text{and} \quad \beta(s(w)) \subset \beta(s(u)),$$

we obtain

$$\alpha(w) \cup \beta(s(w)) \subset N_\delta(\alpha(u) \cup \beta(s(u))),$$

which means that $q(w) \subset N_\delta(q(u))$.

Thus, having in view that $\mu(q(u)) = \mu(q(w))$, we can apply Lemma 1, and we obtain the required inequality $\text{dist}(q(u), q(w)) < \varepsilon$. \square

Notice that n is universal for all points $A, B \in \mu^{-1}(t)$ such that $A \cap B \neq \emptyset$.

We say that a continuum Y is *uniformly continuum-chainable* if for each positive number ε there is an integer $k = k(\varepsilon)$ such that for each pair x, y of points of Y , there are subcontinua A_1, \dots, A_k of Y each of diameter less than ε and such that $x \in A_1$, $y \in A_k$ and $A_i \cap A_j \neq \emptyset$ whenever $|i - j| \leq 1$.

Theorem 1. *If a continuum X is uniformly continuum-chainable, then for any Whitney map μ and any $t \in (0, \mu(X))$, the Whitney level $\mu^{-1}(t)$ is uniformly pathwise connected.*

Proof. Assume that X is a uniformly continuum-chainable continuum. Let μ be a Whitney map for $C(X)$, and let $t \in (0, \mu(X))$ be given.

There exists $\varepsilon' > 0$ such that for $K \in C(X)$ with $\text{diam } K < \varepsilon'$ we have $\mu(K) < t$. Otherwise, we would choose a sequence of elements of $\mu^{-1}([t, \mu(X)])$ convergent to a point, which is impossible for $t > 0$.

For this ε' take an integer k according to the uniform continuum-chainability of X .

Now, fix points A, B of $\mu^{-1}(t)$ and choose $x \in A$ and $y \in B$ arbitrarily.

There are subcontinua A_1, \dots, A_k of X each of diameter less than ε' , and such that $x \in A_1$ and $y \in A_k$ and $A_i \cap A_j \neq \emptyset$ whenever $|i - j| \leq 1$.

We have $\mu(A_i) < t$ for $i = 1, 2, \dots, k$. Thus, for each A_i there is a continuum $B_i \subset X$ such that $A_i \subset B_i$ and $\mu(B_i) = t$.

Let $B_0 = A$ and $B_{k+1} = B$. We have $B_i \cap B_{i+1} \neq \emptyset$, for $i = 0, \dots, k$.

Let $q_i : [0, t] \rightarrow \mu^{-1}(t)$ be a continuous function constructed such as in Lemma 2, and such that $q_i(0) = B_i$ and $q_i(t) = B_{i+1}$, for $i = 0, \dots, k$.

Let $q : [0, (k+1)t] \rightarrow \mu^{-1}(t)$ be such that for any $i = 0, \dots, k$ and $u \in [it, (i+1)t]$ the equality $q(u) = q_i(u - it)$ holds.

Let us observe that q joins A with B .

To show the uniform pathwise connectedness of $\mu^{-1}(t)$, fix $\varepsilon > 0$. For this ε take the integer n according to Lemma 2. By Lemma 2, for the path q the number $(k+1)n$ is that one for which the condition $(**)$ is fulfilled.

Since the numbers n and k do not depend on the choice of points A and B of $\mu^{-1}(t)$, the proof is finished. \square

Now, having Theorem 1, we can answer the questions mentioned at the beginning of the paper.

Theorem 2. *The property of being a uniformly pathwise connected continuum is a Whitney property.*

Proof. Let X be a continuum with the family \mathcal{F} of paths $p:[a,b] \rightarrow X$ satisfying conditions for uniform pathwise connectedness.

By Theorem 1, it suffices to show that X is uniformly continuum-chainable.

Fix $\varepsilon > 0$ and take an integer n according to the uniform pathwise connectedness of X . Let $x, y \in X$ be arbitrarily chosen. There exist $p \in \mathcal{F}$ such that $p(a) = x$ and $p(b) = y$ and points $t_0 = a < t_1 < t_2 < \dots < t_n = b$ of $[a, b]$ such that the diameter of $A_i = p([t_{i-1}, t_i])$ is $\leq \varepsilon$, for $i = 1, 2, \dots, n$, which means that X is uniformly continuum-chainable since $x \in A_1$ and $y \in A_n$. Thus, the proof is finished. \square

We say that a continuum Y is *connected by uniformly short paths* if for each positive number ε there is an integer $k = k(\varepsilon)$ such that for each pair x, y of points of Y , there is a path $p:[a, b] \rightarrow Y$ such that $p(a) = x$, $p(b) = y$, and points $t_0 = a < t_1 < t_2 < \dots < t_k = b$ of $[a, b]$ such that the diameter of $p([t_{i-1}, t_i])$ is $\leq \varepsilon$, for $i = 1, \dots, k$.

This concept was introduced by Bellamy in [1], where he asked whether it was equivalent to the concept of uniform pathwise connectedness. In [3], Holmes answered the question of Bellamy constructing a continuum which is connected by uniformly short paths but not uniformly pathwise connected. Since it is easy to see that a continuum which is connected by uniformly short paths is uniformly continuum-chainable, we have the following.

Theorem 3. *The property of being a uniformly pathwise connected continuum is not a Whitney-reversible property.*

Proof. Take a continuum P which is connected by uniformly short paths but not uniformly pathwise connected. Since P is uniformly continuum-chainable then, by Theorem 1, for each Whitney map $\mu:C(P) \rightarrow [0, \infty)$ and each $t \in (0, \mu(P))$ the Whitney level $\mu^{-1}(t)$ is uniformly pathwise connected. Since the continuum P is not uniformly pathwise connected, the proof is finished. \square

Question. Is it true that a continuum which is uniformly continuum-chainable is connected by uniformly short paths?

Note Added in Proof

The results of this paper were presented at the First International Meeting on Continuum Theory held in Puebla, Mexico, 2000.

Acknowledgements

The author wishes to thank to Professor J. Mioduszewski for the help in completing the work, to Professor J.R. Prais for suggesting the problem, and to Professor P. Krupski for indicating the article [3].

References

- [1] D.P. Bellamy, Short paths in homogeneous continua, *Topology Appl.* 26 (1987) 287–291.
- [2] C.L. Hagopian, L.G. Oversteegen, Continuum chainability without arcs, *Houston J. Math.* 21 (1995) 407–411.
- [3] M.R. Holmes, There is a continuum which is connected by uniformly short paths but not uniformly path connected, *Topology Appl.* 42 (1991) 17–23.
- [4] A. Illanes, S.B. Nadler, *Hyperspaces: Fundamentals and Recent Advances*, Marcel Dekker, New York, 1998.
- [5] H. Kato, A note on continuous mappings and the property of J.L. Kelley, *Proc. Amer. Math. Soc.* 112 (1991) 1143–1148.
- [6] I. Krzemińska, J.R. Prajs, A non- g -contractible uniformly path connected continuum, *Topology Appl.* 91 (1999) 151–158.
- [7] I. Krzemińska, J.R. Prajs, On continua whose hyperspace of subcontinua is σ -locally connected, *Topology Appl.* 96 (1999) 53–61.
- [8] W. Kuperberg, Uniformly pathwise connected continua, in: *Studies in Topology*, Academic Press, 1975, pp. 315–324.
- [9] J. Krasinkiewicz, On the hyperspaces of snake-like and circle-like continua, *Fund. Math.* 83 (1974) 155–164.
- [10] S.B. Nadler Jr, *Hyperspaces of Sets*, Marcel Dekker, 1978.
- [11] S.B. Nadler Jr, *Continuum Theory*, Marcel Dekker, 1992.